

Rec 103/PTO 10 FEB 10

103/584210



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

REC'D 06 OCT 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02078357.7

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:  
Application no.: 02078357.7  
Demande no:

Anmeldetag:  
Date of filing: 13.08.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Akzo Nobel N.V.  
Velperweg 76  
6824 BM Arnhem  
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Replicons of pestviruses that do not express C and or E1 protein and infectious  
viral particles containing same, that can be used in vaccines

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

Title: Replicons of Pestiviruses that do not express C and or E1 protein and infectious viral particles containing same, that can be used in vaccines.

5 The present invention relates to replicons of Pestiviruses that do not express all structural proteins of the virus, infectious viral particles containing said replicons, a method for producing said infectious viral particles and vaccines containing said viral particles.

10 Animals may be protected against pestiviruses by vaccination, however, conventional inactivated or modified live vaccines have disadvantages concerning safety as well as efficacy. Therefore, new types of vaccines should be developed.

Pestiviruses can be divided into two different biotypes, cytopathogenic (cp) and non cytopathogenic (ncp) viruses, respectively. Bovine viral diarrhea virus (BVDV), a member of the genus *Pestivirus* within the family *Flaviviridae* is the causative agent of bovine viral diarrhea, an economically important disease of cattle world-wide. Genetically and structurally closely related virus species are Classical Swine Fever Virus (CSFV) and the ovine Border Disease Virus (BDV). Pestiviruses can induce severe diseases with marked economical losses world wide. The major economic losses caused by BVDV infections are reduced fertility, abortions and the generation of persistently infected calves, which can develop fatal "Mucosal Disease".

20 While cp BVDV induce apoptosis and cell death and express non-structural protein 3 (NS3), inoculation with ncp BVDV leads to persistent infection of cell cultures and NS3-expression is not detectable. The pestivirus genome consists of a single-stranded RNA of positive orientation. The RNA has a length of approximately 12.3 kb and contains one large open reading frame (ORF), which is flanked by non-translated regions (NTR) at both genome ends. The pestiviral ORF is translated into one polypeptide, which is co- and post-translationally processed into 11 (ncp BVDV) or 12 (cp BVDV) mature proteins by viral and cellular proteases. Pestivirus virions consist of four structural proteins, a capsid (C) protein and three glycosylated envelope proteins ( $E^{RNS}$ , E1, E2). BVDV antibodies are directed against  $E^{RNS}$ , E2 and NS3. Neutralizing activity was predominantly demonstrated for E2-specific antibodies.

35 Studies on the replication of pestiviruses have been considerably facilitated by reverse genetic systems and the discovery of autonomously replicating subgenomic RNAs (replicons) (Behrens et al., J.Virol., 72, 2364-272, 1998; Meyers et al., J.Virol., 70, 8606-8613, 1996).

Pestivirus genomes with deletions were first described as defective interfering particles (DI) for BVDV and CSFV (Kupfermann et al., J.Virol., 70, 8175-8181, 1998; Meyers et al., J.Virol., 69, 3683-3689, 1995).

Since then many reports relating to the replication of pestiviruses have been published. *Trans*-complementation of structural proteins of viruses of the family *Flaviviridae* has been reported.

5 Pestivirus self-replicating RNAs are important tools for an understanding of virus replication, assembly and egress. *Trans*-complementation of deleted parts of the genome can be used, for example, for the identification of *trans*-acting elements of a pestiviral genome.

10 The minimal requirements for CSFV replication were investigated, for example, by creating defective CSFV genomes lacking the gene sequences for the structural proteins. It was found that the defective CSFV genomes still replicated and could be packaged into viral particles when introduced in SK-6 cells together with helper A187-CAT RNA (Moser et al., *J.Virol.*, 7787-7794, 1999). An autonomously replicating defective BVDV genome which lacks the genes encoding C, E<sup>ms</sup>, E1, E2, p7 and NS2 had been described (Behrens et al., *J.Virol.*, 72, 2364-2372, 1998).  
15 For the Kunjin virus (KUN) replicons with deletions in the structural regions were packaged into virions using a system where BHK-21 cells were transfected by two consecutive electroporations, first with the mutated KUN replicons, and subsequently with a recombinant Semliki Forest Virus replicon expressing KUN structural proteins (Khromykh et al., *J.Virol.*, 72(7), 5967-5977, 1998). For CSFV it had been recognized  
20 that *trans*-complemented defective virions that contained E<sup>ms</sup> deleted replicons could potentially serve as a basis for vaccines. CSFV, E<sup>RNS</sup>-deleted replicons were generated and *trans*-complemented using a swine kidney cell line (SK6) constitutively expressing CSFV E<sup>RNS</sup>. The resulting virions were able to infect SK6 cells without the production of infectious virus progeny and could be passaged on  
25 E<sup>RNS</sup>-expressing SK6 cells. Pigs were protected against lethal CSFV challenge after immunization with the complemented virions (Widjoatmodjo et al., *J.Virol.*, 74, 2973-2980, 2000).

So far, there are only few reports concerning *trans*-complementation of BVDV replicons. Defects in the coding regions for non-structural proteins NS3, NSNS4a,  
30 NS4B and NS5B were found to be non complementable, whereas defects in the NS5A unit could be complemented in *trans* (Grassmann et al., *J.Virol.*, 75, 7791-7802, 2001). For a mutant BVDV wherein the genes encoding the core, E<sup>ms</sup>, E1 and E2 genes were replaced by the core, E1 and E2 genes of HCV, it was observed that the chimeric viral genome was packaged into virions containing the BVDV structural  
35 proteins contributed by a helper virus (Jae-Hwan Nam, *J.Virol.meth.*, 97, 113-123, 2001). *Trans*-complementation of BVDV E2 and/or p7, using cell lines constitutively expressing BVDV E2 and/ or p7 for complementation have also been described (Harada et al., *J.Virol.*, 74, 9498-9506, 2000).

The present invention provides new Pestiviral RNA genomes (replicons) that are able to replicate, and can be packaged into infectious viral particles in cells that complement the missing protein(s), but do not produce infectious progeny virus. Such replicons can be useful for vaccine purposes.

5 Although it was known in the art that all the genes encoding structural proteins of a Pestivirus can be deleted without affecting the ability to replicate (Behrens et al, *supra*), such a mutant, missing all structural proteins would not be suitable for vaccine purposes. A vaccine mainly aims at invoking an immune response. A  
10 vaccine, on the one hand, should be able to elicit a protective immune response, while, on the other hand, it should of course not invoke the (viral) disease in the inoculated animal or contact animals. The immune response induced is usually mainly directed against the envelop proteins of the virus. But, if a replicon would be  
15 sequences have been deleted, such proteins would not be produced from the replicon and no immune response to these proteins would be obtained.

The present inventors aimed at providing a replicon that would still encode most, preferably all, envelop proteins of the virus, while, on the other hand, it would not be  
20 capable of producing infectious progeny virus.

The present invention provides a replicon of a Pestivirus, preferably the Bovine Viral Diarrhea Virus (BVDV), which expresses all structural proteins of BVDV except for a functional C and/or E1 protein. Preferably at least part of the coding sequence of the  
25 E1 or C protein has been deleted from said replicon.

A replicon is a self-replicating RNA molecule. In this case the RNA molecule is the RNA genome of the BVD virus. Replicons can be constructed from a full length c-DNA clone of the BVD virus, for example from a full-length BVDV cDNA clone like  
30 pA/BVD/ins- (Meyers et al., J. Virol. 70:8606-8613, 1996). The BVDV RNA contains one ORF that is translated into one polyprotein, which is processed into mature proteins by viral and cellular proteases. In constructing mutated replicons care should be taken not to introduce a frameshift within the BVDV ORF. Furthermore, when coding sequences for structural proteins are deleted, the sequences encoding some  
35 N and/or C terminal amino acids may be retained to ensure proper post translational processing of the precursor polypeptide (protease cleavage sites are retained). For example, the capsid (C protein) of BVDV strain NCP7 is encoded by amino acids (AA) 169-270. About 25 amino acids at the C terminus (AA245- 270) function as a signal sequence for the translocation of the E<sup>RNS</sup>E1E2 polyprotein, and as a putative  
40 anchor of the capsid protein. The signal sequence at the C terminus of the capsid protein is essential for a further processing of the polyprotein downstream.

For the C protein a maximum of 32 AA should preferably be retained at the N terminus.

The envelop protein E1 of BVDV strain NCP7 is encoded by amino acids 498 to 692. About 33 amino acids at the C terminus (AA 660 to 692) function as a signal sequence for the translocation of the E2 envelop protein, and as a putative anchor of the E1 envelope protein.

However, other signal sequences or the analogous sequences of other pestivirus species (e.g. CSFV) may be used instead of, for example, the BVDV sequence.

For E1 mutants of BVDV no AA at the N terminus have any function as a signal sequence.

For both E1 and C mutants several ways for construction of replicons are conceivable, that do not express the functional proteins. For example, such mutations may be created by partial deletions whereby the signal sequences are retained, or partial or complete deletions whereby the signal sequences are replaced by analogous signal sequences, or mutations where only base pair exchanges are made.

The present inventors proved for the first time, that both C and E1 structural proteins are essential for the formation of infectious pestiviruses. Furthermore it was shown that deletion of C and E1 does not impact the ability of RNA self-replication.

By using cell lines constitutively expressing Pestiviral structural proteins, Capsid- or E1-proteins can be efficiently *trans*-complemented. The resulting virions are able to infect target cells and to transfer the replicons without generating infectious replication-competent virus progeny. The complemented virions are indistinguishable from wild-type Pestivirus in virus neutralization experiments and can be completely neutralized by, for example in the case of BVDV, a BVDV-specific serum as well as by a BVDV E2-specific serum. Recombinations yielding infectious wild-type BVDV were not detected in any of the complementation experiments.

The complemented viruses may be used for the safe and efficacious immunization against a Pestivirus, such as BVDV.

Infection of cells and animals with the *trans*-complemented viruses results in replicating RNAs expressing all encoded proteins in the infected cells but which do not generate viral particles.

Thus the complemented virions can be used for the immunization against an infection with a pestivirus, such as BVDV, and can protect vaccinated animals from Pestivirus-related disease.

In a preferred embodiment a replicon according to the invention does not encode a functional C protein. This can be achieved by deleting at least part of the coding sequence for the C protein. In case of BVDV, preferably the coding region encoding amino acid positions 201-243 of the C protein is deleted.

In another embodiment of the present invention the replicon does not encode a functional E1 protein. For BVDV, preferably the coding region encoding amino acid positions 498 to 653 of the E1 protein are deleted.

5

It has been found that the C protein, which is the capsid protein of the virus, as well as the E1 protein, is essential for obtaining infectious virus. Mutants that lack the coding sequence for the C or the E1 protein are thus not infectious. The advantage of mutants that do not encode a capsid or E1 protein, but do encode all (envelop) proteins responsible for the immune response (especially E2 and E<sup>ms</sup>) is that the immune response will still include a response against all envelop proteins, since the coding sequences for all envelop proteins is still present.

10

Likewise part of the present invention are infectious viral particles of a pestivirus that contain a replicon according to the invention. Such particles can be incorporated in a vaccine for said Pestivirus, for example BVDV.

15

A method for the production of viral particles according to the invention is likewise part of the invention. Such a method may comprise the following steps:

- Providing cells that are permissive for a particular Pestivirus (for example BVDV) and express a Pestiviral E1 and/or C protein,
- Transfecting said cells with a replicon according to any of claims 1 to 6,
- Culturing transfected cells obtained in step b,
- Harvesting the viral particles from the cell culture.

20

25

Cells that can be used in the production of the viral particles according to the invention should at least be capable of expressing the C protein and/or the E1 protein of a pestivirus, which may be the pestivirus on which the replicon is based or another Pestivirus. In case of BVDV said cells should preferably express the E1 and or C protein of BVDV.

30

In case of BVDV, such cells are preferably of bovine origin, for example bovine kidney cells or diploid bovine esophageal cell line.

The ways by which the coding sequences of a pestivirus such as BVDV can be introduced into said cells are known in the art. For example, the coding sequences for the structural proteins of BVDV (at least C and/or E1) may be introduced into said cells by way of an expression plasmid encoding said proteins or by way of a helper virus. When the viral particles are intended to be used in a vaccine the use of expression plasmids is preferred, to avoid complicated purification methods due to the presence of a helper virus. Suitable expression plasmids are known in the art.

35

40

After transfection of replicon RNA infectious viral particles are produced. The cells can be grown according to methods known in the art until the desired viral titer is obtained. Virus may be harvested from the supernatant of the cell culture or from total cell lysates.

5

Methods used to transfect the cells and the various cloning procedures are known in the art (for example: Sambrook, J & Russell, D.W. Molecular Cloning: A laboratory Manual (Cold Spring Harbor Press, New York, 2001). All permanent cell lines which are suitable for the replication of pestiviruses may be used, preferentially cells of bovine origin. Preferably cells are used that can be easily transfected with DNA. The coding sequences of the structural proteins of most of the pestiviruses contain cryptic splicing sites. Therefore, for most of the BVDV strains, a synthetic open reading frame (synORF) eliminating the splicing sites may be constructed to ensure expression with promoters using the nuclear pathway (e.g. HCMV promoter). However other expression systems (viruses, replicons from other viruses, pestivirus replicons with different deletions) may provide the deleted proteins for trans complementation.

10

15

A vaccine containing infectious viral particles according to the invention together with a pharmaceutically acceptable carrier is likewise part of the present invention.

20

#### BRIEF DESCRIPTION OF THE FIGURES:

##### FIGURE 1.

Schematic representation of the generated constructs. Filled boxes represent the BVDV structural region. Dotted lines show the deleted regions and numbers indicate the nucleotide position in the BVDV full-length RNA. Arrows indicate restriction enzyme sites flanking the deletions. Lines at the left and right ends indicate untranslated regions. N<sup>pro</sup>, autoprotease; C, capsid protein; E<sup>RNS</sup>, E1 and E2, envelope proteins; p7, nonstructural protein; N2 to NS5, nonstructural proteins, 3'-UTR and 5'-UTR, noncoding regions. The scale in kb is given.

25

30

#### EXAMPLES

##### Example 1: Construction and characterization of BVDV replicons.

35

After establishment of an infectious clone for BVDV CP7, a variant of the construct with a deletion of a 27 nucleotide insertion was generated that allowed recovery of ncp BVDV (Meyers et al., J. Virol., 70, 8606-8613, 1996). Based on this construct termed pA/BVDV/Ins-, replicons were constructed by deletion of sequences encoding C and E1 (Fig. 1) without introducing a frameshift within the BVDV ORF. Mutant BVDV clones shown in Fig. 1 were constructed on the basis of the full-length cDNA

40



clone pA/BVDV/ins-. Location of restriction sites are indicated by superscript numbers corresponding to their cleavage positions in the BVDV/CP7 genome (Fig. 1).

## 5 Plasmid constructs

The NCP7ΔC and NCP7ΔE1 plasmids were constructed by a two step cloning procedure. Two PCR fragments were inserted into plasmid pA/BVDV/ins- digested with XhoI<sup>209</sup> and KpnI<sup>2447,3797</sup> (Tab. 1).

- 10 Plasmids were amplified in *E. coli* TOP10F' cells (Invitrogen). Restriction enzyme digestion and cloning procedures were performed according to standard protocols. Plasmid DNA was purified by QIAGEN Plasmid Mini, Midi or Maxi kits. The primers used for PCR or sequencing (labeled with IRD 800) were custom synthesized (MWG-Biotech). Primer pairs for generation of the respective constructs are summarized in Table 1.

- The resulting replicons contained a deletion of a part of the Capsid region (NCP7ΔC, amino acids position 201 to 243 (Fig. 1) or of the E1 encoding region (NCP7ΔE1, amino acid position 498 to 653 (Fig.1).
- 20 All deletions were confirmed by PCR and nucleotide sequencing of the resulting cDNA clones.

## Polymerase chain reaction (PCR) and sequencing

- For PCR, a PTC-200 thermal cycler (MJ Research, Inc.) was used. DNA based amplification was done by Expand High Fidelity PCR System (Roche Molecular Biochemicals) according to the supplier's protocol. For RT-PCR, total RNA of virus-infected cells was extracted using the TRIZOL reagent (Gibco-Life Technologies). cDNA was amplified from the RNA template by using reverse transcriptase (Promega) and a sequence-specific primer. The synthesized cDNA was amplified with a thermostable Taq Polymerase (Promega) and PCR products were directly sequenced.

- Sequencing was carried out using a Thermo Sequenase Cycle Sequencing Kit (Amersham Biosciences). Nucleotide sequences were read with a LI-COR automatic sequencer (MWG Biotech) and were analyzed using the Wisconsin software package version 9.1 (Genetics Computer Group, USA).

## *In vitro*-transcription and electroporation

- Subsequently, *in vitro*-transcribed full-length replicon RNA was transfected into BVDV-negative PT cells.
- 40 *In vitro*-transcription of linearized full-length cDNA constructs and replicons was performed by T7 RiboMax<sup>TM</sup> Large Scale RNA Production System (Promega)

according to the manufacturers instructions. The amount of RNA was estimated by ethidiumbromide staining after agarose gel electrophoresis. For transfections,  $1 \times 10^7$  PT, PT\_805 or KOP-R cells were detached using a trypsin solution, washed with phosphate buffer saline without  $\text{Ca}^{++}/\text{Mg}^{++}$  (PBS<sup>-</sup>), mixed with 1 to 5  $\mu\text{g}$  of *in vitro*-synthesized RNA and electroporated (two pulses at 850V, 25 $\mu\text{F}$ , 156 $\Omega$ ) using an *Easyject Plus* (EquiBio) transfection unit.

#### Characterization of the replicons

For the detection of BVDV proteins, monoclonal antibodies (mab) *WB210* (anti- $\text{E}^{\text{RNS}}$ , CVL, Weybridge), *WB215* (anti-E2, CVL, Weybridge), *CA3* (anti-E2, Institute for Virology, TiHo Hannover), mab-mix *WB103/105* (anti-NS3, CVL, Weybridge), and *C16* (anti-NS3, Institute for Virology, TiHo, Hannover) were used (Edwards et al., Vet.Microbiol., 29,101-108,1991; Edwards et al., Arch.Virol., 102, 197-206, 1988; Peters et al., Vet.Microbiol., 12, 195-200, 1986).

Transient expression of NS3 could be detected from 24 h post transfection by IF for all replicons. Additionally, also  $\text{E}^{\text{RNS}}$ - and E2-specific immunofluorescence was demonstrated. However, no infectious recombinant BVDV could be recovered, even after serial passages and co-passages using highly susceptible KOP-R cells (Table 2). The intensity of NS3-specific immunostaining of NCP7 $\Delta\text{C}$  and NCP7 $\Delta\text{E1}$ -transfected PT cells was comparable to cells transfected with full-length NCP7 RNA and more than 85% of the cells were positive in IF (Table 2). Transfection of full-length NCP7 RNA into PT cells resulted in virus titers of up to  $10^3$  IU/ml. As expected, electroporation of cells with RNA of the prototype replicon DI9 (Behrens et al., J.Virol., 72, 2364-2372, 1998) did not lead to recovery of infectious BVDV, but a strong immunofluorescence following NS3 staining was observed. RT-PCR of PT cells transfected with replicons or full-length NCP7 demonstrated presence of BVDV-specific RNA of the expected size containing the correct deletions. In summary, deleted BVDV genomes, which were replication-competent but lacked the ability of generating virus progeny, were constructed and could efficiently be transfected into bovine cells.

Table\_1: Construction of BVDV-replicons

replicon	deletion (nt)	deletion (aa)	primer	sequence (5'→3')	position (nt)	sense
ncp7ΔC	969-1095	201-243	cp7-204	AAGCCTCGAGATGCCACGTGG	204-224	+
(KpnI <sup>2447</sup> mutated)			C-969R-KpnI	tctaggtaccTCTCTGACTCCTTAGGTGTTATC	969-947	-
			cp7-1096-Kpn	gagagggtaccCTCAAGAGTCGCGCAAGAAAC	1096-1116	+
			p7-3804R	GTCCTAGGTACCCCTGGGCA	3785-3804	-
ncp7ΔE1	1860-	498-653	cp7-204	AAGCCTCGAGATGCCACGTGG	204-224	+
(KpnI <sup>2447</sup> mutated)	2333		cp7-1859R-KpnI	tcagggtaccGTGCATATGCCCCCAAACCATGTC	1859-1838	-
			cp7-2334-KpnI	caacgggtaccGAATTTGGACCCGCTGCTACAACC	2334-2354	+
			p7-3804R	GTCCTAGGTACCCCTGGGCA	3785-3804	-

### Example 2: Establishment of C-E<sup>RNS</sup>-E1-E2 expressing cells

#### **Establishment of C-E<sup>RNS</sup>-E1-E2 expressing PT cells**

5 In order to do complementation studies with the BVDV replicons, a bovine cell line was established that constitutively expresses BVDV structural proteins.

10 PT cells (RIE11 CCVL), a permanent bovine kidney cell line, was obtained from the collection of cell lines in veterinary medicine at the Federal Research Center of Virus diseases of Animals, Insel Riems (CCLV). PT cells were chosen due to their applicability for DNA/RNA transfection and for generation of constitutively expressing cell lines. Cells were grown in Dulbecco's minimal essential medium (DMEM) supplemented with 10% BVDV-free fetal bovine serum (FBS).

15 The genomic region encoding the structural proteins (C-E<sup>RNS</sup>-E1-E2) of BVDV strain PT810 (Wolfmeier et al., Arch.Virol., 142, 2049-2057, 1997) was cloned as a chemically synthesized synthetic open reading frame (Syn-ORF; kindly provided by Tobias Schlapp, Bayer AG, Leverkusen). It consisted of 2694 nucleotides extending from nucleotide 890 to 3584 of the nucleotide sequence of BVDV strain NADL (Collett et al., Virology, 165, 200-208, 1988) and was inserted into the pcDNA3.1 expression plasmid (Invitrogen) using restriction sites *KpnI* and *NotI*. The nucleotide  
20 sequence of Syn-ORF had been changed to remove splice sites (Schmitt et al., J.Gen.Virol., 80, 2839-2848, 1999), but retained the original amino acid sequence of BVDV strain PT810. Additionally, the first codon of Syn-ORF was changed to a methionin to allow expression of the polyprotein under the control of the *HCMV* Immediate early promotor present in pcDNA3.1, and a stop codon was inserted  
25 behind the last codon.

The resulting construct pcDNA\_C-E2 (1 µg) was used to transfect PT cells with the SUPERFECT reagent (Qiagen). At 2 days after transfection, cell culture medium was changed to DMEM supplemented with 10% bovine serum and 1mg of G418 (Gibco,  
30 Life Technologies) per ml. G418-resistant colonies were isolated, and replated several times.

After G418 selection and passaging, resistant cell clones were tested for BVDV E<sup>RNS</sup> and E2 expression using mabs WB210 and WB215, respectively. Positive cells were cloned, passaged under G418 selection and tested again for E<sup>RNS</sup> and E2  
35 expression. In the IF analyses, G418-resistant cell clone PT\_805 contained more than 90% E<sup>RNS</sup> and E2 expressing cells. During further passaging of PT\_805 cells for 6 months in the presence of G418, the portion of E<sup>RNS</sup> and E2 expressing cells remained stable. Immunoprecipitation experiments with E<sup>RNS</sup>- and E2-specific mabs showed a protein pattern identical to that of BVDV infected PT cells, where  
40 glycoprotein E1 is co-precipitated with the E2-specific mab (data not shown). From these data and those obtained by nucleotide sequencing it was concluded that all

BVDV structural proteins were constitutively expressed in PT\_805 cells. The cell line was used for *trans*-complementation experiments and for the generation of infectious virions as well as for packaging and transducing deleted replicons.

#### 5 Recovery and characterization of *trans*-complemented recombinant BVDV

For *trans*-complementation experiments, *in vitro*-transcribed BVDV replicon RNA or full-length NCP7 RNA was transfected into monolayers of PT\_805 cells by electroporation. Cell culture supernatants were collected 24h, 48 h and 72h *post transfection* (p.t.), clarified by centrifugation (10,000 x g, 5 min) and titrated using highly susceptible KOP-R cells. KOP-R cells (RIE244, CCVL) are cells of a diploid bovine esophageal cell line, obtained from the collection of cell lines in veterinary medicine at the Federal Research Center of Virus diseases of Animals, Insel Riems (CCLV). KOP-R cells were selected due to their susceptibility to BVDV infection and their suitability for BVDV propagation. Cells were grown as described above for the PT cells. As a negative control for replication, replication incompetent controls were used (deletion mutants NCPΔNS5 and NCP7Δ3'nteraatl).

At the day of collection, replication of BVDV was monitored by IF staining.

Virus titers of the complemented viruses were determined as infectious units (IU). Cell culture supernatants were titrated in triplicate in log<sub>10</sub> steps and 1 ml was inoculated onto KOP-R cells seeded in 24-well plates. After 12 h of incubation at 37°C, cells were washed, detached with a trypsin solution, and counted. An aliquot of the cells was stained by IF using a BVDV NS3-specific mab and analyzed by flow cytometry. Infectious units (IU) were calculated according to the formula:

number of cells in the well x % IF-positive cells x dilution factor = titer in IU/ml

For calculations, only wells with >5% and <30% IF-positive cells were considered.

For all replicons and full-length NCP7, NS3 staining following RNA transfection led to immunofluorescence patterns similar to that obtained with PT cells. Between 80% and 95% of transfected cells were reactive with NS3-specific antibodies. Infectious recombinant BVDV was detected in supernatants of NCP7ΔC, NCP7ΔE1 and NCP7-transfected PT 805 cells as shown by infection of KOP-R cells. Virus titers of the resulting complemented viruses NCP7ΔC<sub>trans</sub> and NCP7ΔE1<sub>trans</sub> varied between 10<sup>4.0</sup> to 10<sup>6.8</sup> IU/ml at 24 h p.t. (Table 2). After transfection of full-length NCP7 RNA into complementing PT\_805 cells, virus titers of up to 10<sup>7.3</sup> IU/ml could be detected at 24 p.t. (Table 2). Therefore, titers of *trans*-complemented viruses from supernatants collected at 24 h p.t. were approximately 10000-fold higher than virus titers obtained after transfection of full-length NCP7 RNA into non complementing PT cells (Table 2). In addition, no complemented BVDV was detected after passaging of PT\_805 cell culture supernatants transfected with DI9 replicon RNAs (Behrens et al., J.Virol., 72, 2364-2372, 1998) at all times p.t. (Table 2).

Using IF of KOP-R cells infected with NCP7ΔC\_trans and NCP7ΔE1\_trans, expression of E<sup>RNS</sup>, E2 and NS3 could be detected (Table 2). Expression pattern indicated that supernatants of transfected PT\_805 cells contained infectious virus that can infect and replicate in KOP-R cells. Titration experiments and subsequent IF staining demonstrated that no cell-to-cell spread occurred, and that neither NCP7ΔC\_trans nor NCP7ΔE1\_trans were able to form NS3-positive plaques or secondary infections in these cells (Table 2). At higher dilutions, only single cells or small foci of cells were positive by IF-staining for NS3. Control experiments with BVDV NCP7 on KOP-R cells resulted in antigen positive virus plaques as demonstrated for E<sup>RNS</sup>, NS3 and E2.

The infection of KOP-R cells with *trans*-complemented viruses could be blocked with BVDV-neutralizing antisera (S-BVD\_pos, S-BVD\_E2) and no infectivity was detected after inoculation of KOP-R cells (100% neutralization).. For neutralization of BVDV, 100 µl of virus suspension (10<sup>4</sup> TCID<sub>50</sub>/ml) was mixed in a 24-well plate with 50 µl of undiluted serum and incubated at 37°C. After 2 h, KOP-R cells (2 x 10<sup>4</sup>) were transferred to the mixture and incubated for 3 days at 37°C. Subsequently, the monolayer was stained for BVDV-NS3 by indirect immunofluorescence (IF) (Grummer et al., J.Gen.Virol., 82, 2597-2605, 2001).

(S-BVD\_pos) was a pooled anti-BVDV serum collected from cattle experimentally infected with BVDV strain PT810 (Beer et al., I.Vet.Microbiol., 77, 195-208, 2000). The homologous neutralization titer of this serum was 1:4,096. For E2-specific neutralization, serum from sheep, immunized with an E2-expressing modified Vaccinia Virus Ankara (MVA) was used (S-BVD\_E2; Beer et al., unpublished data).

In a control reaction, incubation with a BVDV antibody-free cattle serum (S-BVD\_neg) did not result in virus neutralization (Table 3).

TABLE 2. Replication and trans-complementation of BVDV NCP7 RNAs with *in-frame* deletions (replicons).

construct	replication <sup>a</sup> after RNA transfection in PT cells (24 h)	1st passage <sup>b</sup> (24 h, 48 h, 72 h) on KOP-R-cells	Titer of virus progeny <sup>c</sup> 24 h after transfection of complementing PT_805 cells (IU/ml)	4th passage <sup>d</sup> of virus progeny on KOP-R cells
NCP7	++ (P) <sup>e</sup>	+ (10 <sup>3</sup> IU/ml)	10 <sup>7.3</sup> (P)	+
NCP7ΔC	++ <sup>f</sup>	Ø <sup>f</sup>	> 10 <sup>5.0</sup>	Ø
NCP7ΔE1	++	Ø	> 10 <sup>5.0</sup>	Ø
DI9	+++	Ø	< 10 <sup>0</sup>	nd

<sup>a</sup> PT cells were transfected with *in vitro*-transcribed RNA and IF-stained for NS3-expression after 24 h.

<sup>b</sup> KOP-R cells were inoculated with undiluted supernatants 24 h, 48 h and 72 h after transfection of PT cells (1 ml per 10<sup>5</sup> cells).  
KOP-R cells were stained with NS3-specific mabs 5 days post inoculation.

<sup>c</sup> PT\_805 cells were transfected with *in vitro*-transcribed RNA and the supernatants were titrated after 24 h using KOP-R-cells.

<sup>d</sup> KOP-R cells were inoculated using supernatants of transfected PT\_805 cells with a titer > 10<sup>6</sup> IU/ml (1 ml per 10<sup>5</sup> cells).

<sup>e</sup> P Indicates the detection of virus plaques after NS3-staining (> 10 cells).

<sup>f</sup> Ø = no BVDV-specific immunofluorescence; + = weak positive NS3-IF-signal and < 50% IF-positive cells, ++ = positive NS3-IF-signal and > 50% IF-positive cells, +++ = strong positive NS3-IF-signal and > 50% IF-positive cells.

nd = not done

TABLE 3. Neutralization of wild-type and complemented BVD viruses by BVDV-specific antisera

BVDV <sup>a</sup>	IF analysis 3 days post inoculation of the neutralization mix		
	S-BVD_pos <sup>b</sup>	S-BVD_E2 <sup>c</sup>	S-BVD_neg <sup>d</sup>
NCP7	Ø	Ø	+ <sup>e</sup>
NCP7AC_trans	Ø	Ø	+
NCP7AE1_trans	Ø	Ø	+

<sup>a</sup> Complemented viruses were adjusted to a titer of 10<sup>4</sup> IU/ml.<sup>b</sup> Cattle serum prepared against BVDV type 1.<sup>c</sup> Sheep serum specifically directed against BVDV type 1 E2.<sup>d</sup> BVDV antibody free cattle serum.<sup>e</sup> Ø = no IF-positive cells, + = more than 100 IF-positive cells/well.



**CLAIMS:**

1. A replicon of a pestivirus which is incapable of expressing one or more structural proteins of the virus, characterized in that said replicon expresses all structural proteins of a pestivirus except for a functional C and/or E1 protein.
2. A replicon according to claim 1, characterized in that at least part of the coding sequence of the E1 or C protein has been deleted from said replicon.
3. A replicon according to claim 1 or 2, characterized in that said replicon does not encode a functional C protein.
4. A replicon according to claim 1 or 2, characterized in that said replicon is of the Bovine Viral Diarrhea Virus (BVDV).
5. A replicon according to claim 4, characterized in that the coding region encoding amino acid positions 201-243 of the C protein have been deleted.
6. A replicon according to claim 1 or 2, characterized in that said replicon does not encode a functional E1 protein.
7. A replicon according to claim 4, characterized in that the coding region encoding amino acid positions 498 to 653 of the E1 protein have been deleted.
8. Infectious viral particle of Pestivirus, characterized in that it contains a replicon according to any of claims 1-7.
9. A method for the production of viral particles of a Pestivirus according to claim 8, characterized in that said method comprises the following steps:
  - a. Providing cells that are permissive for the Pestivirus and express Pestiviral E1 and/or C protein,
  - b. Transfecting said cells with in-vitro transcribed RNA of a replicon according to any of claims 1 to 7,
  - c. Culturing transfected cells obtained in step b,
  - d. Harvesting the viral particles from the cultured cells.
10. A method according to claim 9, characterized in that said pestivirus is BVDV.
11. A method according to claim 10 or 11, characterized in that said cells express the E1 and/or C protein of BVDV.
12. A vaccine containing infectious viral particles according to claim 8 and a pharmaceutically acceptable carrier.

## ABSTRACT

The present invention provides new Pestiviral RNA genomes (replicons) that are able to replicate, and can be packaged into infectious viral particles in cells that complement the missing protein(s), but do not produce infectious progeny virus. Such replicons can be useful for vaccine purposes.

The replicons encode most, preferably all, envelop proteins of the virus, while, on the other hand, it would not be capable of producing infectious progeny virus.

The present invention provides a Pestiviral replicon, preferably from the Bovine Viral Diarrhea Virus (BVDV), which expresses all structural proteins except for a functional C or E1 protein. Preferably at least part of the coding sequence of the E1 or C protein has been deleted from said replicon.


The present inventors proved for the first time, that both C and E1 structural proteins are essential for the formation of infectious pestiviruses. Furthermore it was shown that deletion of C and E1 does not impact the ability of RNA self-replication.

By using cell lines constitutively expressing pestiviral structural proteins, Capsid- or E1-proteins can be efficiently *trans*-complemented. The resulting virions are able to infect bovine target cells and to transfer the replicons without generating replication-competent virus progeny. In other words, no infectious progeny virus is produced. The complemented virions are indistinguishable from wild-type Pestivirus in virus neutralization experiments. Recombinations yielding infectious wild-type virus were not detected in any of the complementation experiments.

The complemented viruses may be used for the safe and efficacious immunization against BVDV.

Figures 1/1

FIGURE 1



PCT Application

**EP0309031**

